AD	

Award Number: DAMD17-00-1-0036

TITLE: The Role of aVb6-mediated latent TGF-B1 Activation in Prostate Cancer

PRINCIPAL INVESTIGATOR: John S. Munger, M.D.

CONTRACTING ORGANIZATION: New York University School of Medicine

New York, New York 10016

REPORT DATE: January 2002

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command

Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;

Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

transforming growth factor-B, integrin, TGFB activation, PTEN,

OF THIS PAGE

18. SECURITY CLASSIFICATION

Unclassified

phosphatase, cell proliferation, epithelium

Form Approved OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE	3. REPORT TYPE AND	DATES COVERE	D
	January 2002	Annual (1 Jan		
4. TITLE AND SUBTITLE			5. FUNDING N	
The Role of aVb6-mediate	d latent TGF-B1 Activ	ation in	DAMD17-00-	-1-0036
Prostate Cancer				
6. AUTHOR(S)				
John S. Munger, M.D.				
7 DEDECORAING ODGANIZATION NAS	ME(S) AND ADDRESS/ES)		9 DEDECIDATING	G ORGANIZATION
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)			REPORT NUMBER	
New York University School of Medicine New York, New York 10016				
New Tork, New Tork Tool	.0			
E saut, mungai04@mad muu adu				
E-Mail: mungej01@med.nyu.edu			<u> </u>	
9. SPONSORING / MONITORING AGE	NCV NAME(S) AND ADDRESS/ES	3)	10 SPONSORII	NG / MONITORING
9. SPONSORING / MONITORING AGE	MOT MAINE(S) AND ADDRESS(EC	<i>'</i> 1	1	EPORT NUMBER
U.S. Army Medical Research and M	Materiel Command			
Fort Detrick, Maryland 21702-5013			1	
· •			Ĭ	
			ĺ	
11. SUPPLEMENTARY NOTES	· · · · · · · · · · · · · · · · · · ·		J	
THE GOLD ELIMENTARY WOLLD				
12a. DISTRIBUTION / AVAILABILITY S				12b. DISTRIBUTION CODE
Approved for Public Rele	ase; Distribution Unl	imited		
13. ABSTRACT (Maximum 200 Words	s)			
Abundant evidence sugges		on of TGF-B by r	orostate car	ncer cells enhances
their ability to grow as				
results from a noncovale				
associated peptide, LAP				
present. Our lab discove				
binds to an RGD sequence				
cells. We hypothesize th				
prostate epithelial pro				
acts in concert with ove				
and promote growth of the tumor. In this work, we are testing whether the ß6 integrin subunit is regulated by androgen, whether it is overexpressed in human prostate cancer,				
and whether it affects of				
results to date indicate				
fashion after castration				
involution is affected h				
develop prostate cancer			· ·	producing mice that
develop prostate cancer	(IRAHI) CHAC AISO IAC	no gene expre	331011.	
14. SUBJECT TERMS				15. NUMBER OF PAGES

OF REPORT

17. SECURITY CLASSIFICATION

Unclassified

20. LIMITATION OF ABSTRACT

Unlimited

16. PRICE CODE

19. SECURITY CLASSIFICATION

Unclassified

OF ABSTRACT

Table of Contents

Cover	1
SF 298	
Introduction	3
Body	4
Key Research Accomplishments	5
Reportable Outcomes	6
Conclusions	7
References	
Appendices	N/A

INTRODUCTION

The subject of this work is a system for the activation of latent TGFB in the prostate. The system consists of the $\alpha V\beta$ 6 integrin expressed on epithelial cells. Our previous work showed that this integrin can bind to an integrin recognition site (arg-gly-asp) on latent TGF\$1 and effect its activation [1]. TGFB is known to be important for regulating the growth and differentiation of various epithelia, and also to be important in cancer growth. Little is currently known about this system in the prostate: eg, what cells express $\alpha V \beta 6$, how expression of the integrin is regulated, and if and when this system regulates prostate epithelial growth via production of active TGFB1. The purpose of the work is to demonstrate whether or not this system plays a role in prostate cancer. The scope of the work involves cell line and mouse experiments (to gauge the normal expression and regulation of $\alpha V \beta 6$ in the prostate), and evaluation of human prostate cancer tissue and an in vivo mouse prostate cancer model (to address the question, does $lpha extsf{V}eta 6 ext{-mediated}$ activation of TGF&1 promote prostate cancer growth?).

BODY

The second 12 months of the project addressed tasks 1 (determine &6 expression in normal prostate and prostate cancer cells), 2 (determine the effect of androgen on &6 expression), 3 (effect of &6 on prostate epithlial proliferation) and 5 (effect of &6 on cancer in a mouse model of prostate cancer).

We have castrated control and ß6 KO mice and sacrificed them at multiple time points. We have also BrDU-labeled control and KO mice. We now have sufficient samples to analyze and are systematically analyzing the samples. We are doing TUNEL staining of the involuting prostate specimens to see if ß6 expression alters the rate of apoptosis. Preliminary results suggest that at the day 3 time point there is less apoptosis in the KO mice. The only remaining samples to generate are exvoluting prostates in normal and KO mice (castrated mice are given DHT pellets to induce regrowth of the prostate).

We have obtained TRAMP mice and are currently crossing them with the ß6 KO mice. Once we generate combined ß6KO/TRAMP (estimate: 2-3 months) mice we will assess their tumor growth and metastasis.

We have looked at ß6 expression in human prostate cancer tissue. In normal areas, ß6 is expressed in basaol cells. In the cancer areas, we see no ß6 expression except in occasional intensely stained cells. The tumors are of intermediate grades and we are trying to get samples of higher grade tumors as well as of metastases to see if s6 expression is a marker of more aggressive cancer.

KEY RESEARCH ACCOMPLISHMENTS

- \$6 is expressed in basal cells of human prostate. This may be important for growth control of these cells (via TGf\$\beta\$ activation) and might be involved in growth control of prostate stem cells, which may exist in this compartment.
- Human prostate cancer, at least of intermediate grade, does not express s6 except for rare positive cells. This may be because these lower grade tumors are still growth-inhibited by TGFB, or because they are of luminal origin.

REPORTABLE OUTCOMES

The human prostate data are reportable.

CONCLUSIONS

These experiments are the first to describe ß6 expression in human prostate cancer, and in normal prostate. The remaining work done has not yet been fully analyzed but should give insights into the role of ß6 integrin on prostate epithelial cell proliferation during androgen withdrawal and replacement.

<u>"So what."</u> There are many reports in the literature relating cancer outcomes and tumor cell behavior to increased expression of TGFß by tumor cells. However, to my knowledge there has never been an analysis of the role of a TGFß activator in tumor cells. Yet, our results with $\alpha V\beta 6$ and lung fibrosis [1] point out the critical role that a TGFß activator can play in a TGFß-dependent process. If a specific TGFß activator can be identified as important in a cancer, this knowledge might be important for determining prognosis and for developing therapies in which the activator is a target.

REFERENCES

- 1. Munger, J.S., et al., The integrin $\alpha v \beta 6$ binds and activates latent TGF $\beta 1$: a mechanism for regulating pulmonary inflammation and fibrosis. Cell, 1999. **96**: p. 319-328.
- 2. Salm, S., et al., Generation of active TGF-B by prostatic cell cocultures using novel basal and luminal prostatic epithelial cell lines. Journal of Cellular Physiology, 2000. 184: p. 70-79.